



Review

New insights into physiology and metabolism of *Propionibacterium freudenreichii*

Anne Thierry ^{a,b,*}, Stéphanie-Marie Deutsch ^{a,b}, Hélène Falentin ^{a,b}, Marion Dalmasso ^{a,b},
Fabien J. Cousin ^{a,b}, Gwenaël Jan ^{a,b}

^a INRA, UMR1253 Science et Technologie du Lait et de l'Œuf, 65 rue de Saint Brieuc, 35000 Rennes, France

^b AGROCAMPUS OUEST, UMR1253 Science et Technologie du Lait et de l'Œuf, 65 rue de Saint Brieuc, 35000 Rennes, France

ARTICLE INFO

Available online 8 May 2011

Keywords:

Propionibacteria
Metabolism
Survival
Swiss cheese
Probiotic

ABSTRACT

Dairy propionibacteria are Actinobacteria, mainly isolated from dairy environments. *Propionibacterium freudenreichii* has been used for a long time as a ripening culture in Swiss-type cheese manufacture, and is more and more considered for its potent probiotic effects. This review summarises the knowledge on the main *P. freudenreichii* pathways and the main features explaining its hardiness, and focuses on recent advances concerning its applications as a cheese ripening agent and as a probiotic for human health. Propionibacteria have a peculiar metabolism, characterised by the formation of propionic acid as main fermentation end-product. They have few nutritional requirements and are able to use a variety of carbon substrates. From the sequence of *P. freudenreichii* CIRM-BIA1^T genome, many pathways were reconstituted, including the Wood–Werkman cycle, enzymes of the respiratory chain, synthesis pathways for all amino acids and many vitamins including vitamin B₁₂. *P. freudenreichii* displays features allowing its long-term survival. It accumulates inorganic polyphosphate (polyP) as energy reserve, carbon storage compounds (glycogen), and compatible solutes such as trehalose. In cheese, *P. freudenreichii* plays an essential role in the production of a variety of flavour compounds, including not only propionic acid, but also free fatty acids released via lipolysis of milk glycerides and methyl-butanoic acids resulting from amino acid degradation. *P. freudenreichii* can exert health-promoting activities, such as a bifidogenic effect in the human gut and promising immunomodulatory effects. Many *P. freudenreichii* properties involved in adaptation, cheese ripening, bio-preservation and probiotic effects are highly strain-dependent. The elucidation of the molecular mechanisms involved is now facilitated by the availability of genome sequence and molecular tools. It will help in the selection of the most appropriate strain for each application.

© 2011 Elsevier B.V. All rights reserved.

Contents

1.	Introduction	20
2.	General features of propionibacteria	20
2.1.	Propionibacteria taxonomy	20
2.2.	Genetic diversity of <i>P. freudenreichii</i>	20
2.3.	Propionibacteria ecology	20
2.4.	Safety aspects	21
3.	<i>P. freudenreichii</i> , a hardy bacterium with original metabolic pathways	21
3.1.	Propionic fermentation via the Wood Werkman cycle: a high energy yield	21
3.2.	Respiration	22
3.3.	Vitamin and porphyrin synthesis	22
3.4.	Production of exopolysaccharides	22
3.5.	Features allowing long-term survival	22
4.	Recent advances in knowledge on <i>P. freudenreichii</i>	23
4.1.	Evolution of published data about dairy propionibacteria	23
4.2.	Role in aroma formation in cheese	23

* Corresponding author at: UMR1253 Science et Technologie du Lait et de l'Œuf, 65 rue de Saint Brieuc, 35000 Rennes, France. Tel.: +33 2 2348 5337; fax: +33 2 2348 5350.
E-mail address: anne.thierry@rennes.inra.fr (A. Thierry).

4.3. Probiotic properties of <i>P. freudenreichii</i>	24
4.4. Protective cultures	25
5. Concluding remarks	25
References	25

1. Introduction

Propionibacteria are Gram positive, high G + C%, mesophilic, aerotolerant, pleomorphic rods. They have a peculiar metabolism leading to the formation of propionic acid as main end-product of fermentation. They display low nutritional requirements and are characterised by their long-term survival in different environments.

P. freudenreichii was first described more than one century ago in Swiss Emmental cheese by Orla Jensen and von Freudenreich (von Freudenreich and Orla-Jensen, 1906), who showed the relationship between the presence of these bacteria producing propionic acid and the formation of the characteristic round holes (eyes) in cheese.

P. freudenreichii is widely used as a ripening culture in the manufacture of Swiss-type cheeses (Dorner, 1939; Langsrud and Reinbold, 1973a; Steffen et al., 1993). It is also known for its production of vitamin B₁₂ and propionic acid, and has properties as protective cultures in food and feed (Thierry et al., 2011). Moreover *P. freudenreichii* is also more and more studied for its probiotic properties (Cousin et al., 2010).

This review summarises the knowledge on the main metabolic pathways of *P. freudenreichii* and the main characteristics explaining its hardiness, and focuses on recent advances concerning its applications as a cheese ripening agent and as a probiotic for human health.

2. General features of propionibacteria

2.1. Propionibacteria taxonomy

Propionibacteria belong to the Actinobacteria class, comprising high G + C content Gram-positive bacteria (Stackebrandt et al., 1997). Actinobacteria exhibit a wide range of morphologies, lifestyles, physiological and metabolic properties and colonise various niches including soil (Ventura et al., 2007). The *Propionibacterium* genus currently comprises 12 species (<http://www.bacterio.cict.fr>). Two distinct groups are distinguished on the basis on their natural biotopes: “dairy” (or “classical”) and “cutaneous” propionibacteria (Cummins and Johnson, 1986). Dairy propionibacteria have been traditionally isolated from milk and dairy products. Four typical dairy species were early described: *P. freudenreichii*, *P. acidipropionici*, *P. jensenii* and *P. thoenii*. Cutaneous propionibacteria are commensal of mammals including humans, the most studied species within this group being *P. acnes*, involved in acne and in post surgery infections. Four propionibacteria species, isolated from other biotopes than dairy products and human skin, were more recently described: *P. cyclohexanicum* isolated from spoiled orange juice (Kusano et al., 1997), *P. microaerophilum* from olive mill wastewater (Koussemon et al., 2001), *P. australiense* from granulomatous bovine lesions (Bernard et al., 2002), and *P. acidifaciens*, isolated from human mouth (Downes and Wade, 2009). The analysis of 16S rRNA gene sequences showed that *P. cyclohexanicum*, *P. acidifaciens* and *P. australiense* are phylogenetically related to *P. freudenreichii*, whereas *P. microaerophilum* is related to *P. acidipropionici* (Downes and Wade, 2009).

2.2. Genetic diversity of *P. freudenreichii*

The genome size of *P. freudenreichii* ranges around 2.6 Mb and its G + C content is 67%. The first genome of a *P. freudenreichii* strain (CIRM-BIA1^T)

has recently been sequenced (Falentin et al., 2010a). Furthermore, molecular tools have newly been developed to over pass *P. freudenreichii*'s poor efficiency of transformation (from 10³ to 10⁸ colony-forming units (cfu)/μg of DNA), thus allowing the exploration of the role of specific genes of interest (Thierry et al., in 2011; Van Lwijk et al., 2002).

Data on genomic biodiversity within *P. freudenreichii* species are scarce. Fingerprinting methods such as Pulsed-Field Gel Electrophoresis and Randomly Amplified Polymorphic DNA-PCR are available to characterise *P. freudenreichii* at the strain level (Meile et al., 2008). Multi-locus sequence typing (MLST) based on the sequence analysis of internal fragments of 7 genes has recently been developed for *P. freudenreichii*. It was applied to 113 strains of different phenotypes and origins (Dalmaso et al., 2011). MLST resolved 46 sequence types grouping each 1 to 11 strains. This study showed that the *P. freudenreichii* core genome possesses a low level of nucleotide polymorphism and that recombination played a significant role in the distribution of this polymorphism among isolates (Dalmaso et al., 2011).

P. freudenreichii was in the past divided into two subspecies on the basis of lactose fermentation and nitrate reductase activity. *P. freudenreichii* subsp. *freudenreichii* is unable to ferment lactose and shows nitrate reductase activity whereas *P. freudenreichii* subsp. *shermanii* exhibits the opposite properties (Cummins and Johnson, 1986). However, the existence of *P. freudenreichii* strains harbouring the two other possible phenotypes (lactose and nitrate reductase positive, and lactose and nitrate reductase negative) has also been mentioned (Dalmaso et al., 2011; de Carvalho et al., 1994; Moore and Holdeman, 1986; Vorobjeva, 1999). In the same way, annotation of the complete *P. freudenreichii* CIRM-BIA1^T genome revealed that the *lacZ* gene responsible for betagalactosidase activity is harboured by a transposon-like mobile element and that the locus responsible for nitrate reduction activity harbours some pseudogenes (Falentin et al., 2010a). Moreover, the phylogenetic analysis of *P. freudenreichii* population using MLST did not fit with the distinction of subspecies (Dalmaso et al., 2011). All these results suggest that there seems to be no justification for separation at the subspecies level according to these two phenotypic criteria.

2.3. Propionibacteria ecology

Strains of dairy origin constitute the main part (80 to 90%) of the *P. freudenreichii* strains available in the international collections (for example, in the Belgian Co-ordinated Collections of Micro-organisms (BCCMTM)/LMG collection and in the CIRM-BIA (Centre International de Ressources Microbiennes – Bactéries d'Intérêt Alimentaire, INRA, Rennes, France), and in other large laboratory collections (Fessler, 1997). However, a few *P. freudenreichii* strains have also been isolated from other biotopes, like hay and straw. Its known habitats are related to cattle environment but would require further investigations, since no systematic search has been made for propionibacteria in habitats distinct from dairy and cattle environments.

In most cheeses, *P. freudenreichii* and other dairy propionibacteria are present at low population density. In Emmental and similar cheeses, however, *P. freudenreichii* reaches a high population density, with counts over 10⁹ cfu/g of cheese during the whole ripening time. The high thermotolerance of *P. freudenreichii*, compared to the other dairy species, would be responsible for the prevalence of this species

in Emmental-type cheeses in which the curd is heated at 50–55 °C for about 30 min.

2.4. Safety aspects

P. freudenreichii has a long history of safe use in human diet and animal feed. It has been granted the Generally Recognized As Safe (GRAS) status from the US Food and Drug Administration (Mogensen et al., 2002). *P. freudenreichii* also belongs, as well as *P. acidipropionici*, to the list of agents recommended for Qualified Presumption of Safety (QPS) by the European Food Safety Authority (EFSA, 2009). QPS is a generic risk assessment approach introduced by EFSA for harmonising the assessment of notified biological agents intended to be introduced into the food chain (Leuschner et al., 2010).

The antibioresistance of the strains intended to be used in food and feed should be investigated. Data about the antibioresistance of *P. freudenreichii* and other dairy propionibacteria species are very scarce. Dairy propionibacteria have intrinsic or “natural” resistance to several antibiotics, including sulfonamides, oxacillin, aminoglycosides, 1st and 2nd generation quinolones, colistin, metronidazole and fosfomycin (Cummins and Johnson, 1992; Meile et al., 2008; Suomalainen et al., 2008). No mobile DNA encoding antibioresistance was detected in dairy propionibacteria. The studies on antibiotic resistance of propionibacteria mainly focused on cutaneous species of clinical importance, especially on *P. acnes* isolates (Nord and Oprica, 2006). Minimal Inhibitory Concentration (MIC), defined as the lowest concentration of the antimicrobial that inhibits bacterial growth, have been recently proposed in a Technical Guidance of EFSA for some bacteria used in animal feed including *Propionibacterium*, but were mainly based on data extrapolated from cutaneous propionibacteria studies (EFSA, 2008).

3. *P. freudenreichii*, a hardy bacterium with original metabolic pathways

Propionibacteria exhibit some peculiar metabolic pathways that have been thoroughly investigated (see reviews (Hettinga and Reinbold, 1972; Piveteau, 1999; Wood, 1981)). The metabolism of propionibacteria is very complex, because several interconnected

pathways are used simultaneously. Propionibacteria are anaerobic but are also aerotolerant. They have low nutritional requirements and are able to fit, survive and remain active in various environments. This section summarises the knowledge on the main pathways of *P. freudenreichii* and the main features explaining its hardiness.

3.1. Propionic fermentation via the Wood–Werkman cycle: a high energy yield

P. freudenreichii ferments a variety of substrates, including carbohydrates, polyols like glycerol and erythritol and adonitol, and organic acids like lactic and gluconic acids (Cummins and Johnson, 1986). The main fermentation products are propionic, acetic, succinic acids and CO₂. Substrates are first oxidised to pyruvate *via* glycolysis or *via* the pentose phosphate pathway, generating ATP and reduced co-enzymes. Pyruvate is oxidised to acetate and CO₂ or reduced to propionate.

The reduction of pyruvate to propionate occurs *via* a specific cycle referred to as transcarboxylase cycle or Wood–Werkman (Crow, 1987; Houwen et al., 1991) (Fig. 1). This pathway uses NADH formed during glycolysis and pyruvate oxidation, and produces extra-ATP. All the reactions of the transcarboxylase cycle are reversible. One of the key reactions is a transcarboxylation reaction transferring a carboxyl group from methylmalonyl-CoA to pyruvate to form oxaloacetate and propionyl-CoA, and occurring without ATP consumption. The enzyme involved is a complex, biotin-dependent carboxytransferase, composed of three subunits of known structure. Oxaloacetate is then converted to succinate, with the reactions catalysed by the enzymes of the citric acid cycle. The reduction of fumarate to succinate is catalysed by a membrane-bound succinate dehydrogenase involving anaerobic electron transport by cytochrome *b* and generation of ATP. Succinate is then converted to succinyl-CoA, concomitantly with the conversion of propionyl-CoA to propionate, by a CoAtransferase. Succinyl-CoA is finally isomerised to methylmalonyl-CoA in a reaction catalysed by methylmalonyl-CoA mutase (EC 5.4.99.2). This coenzyme B₁₂-dependent enzyme is another original enzyme of propionibacteria. In summary, the pathway of propionate production in propionibacteria is unique, functions as a cyclic process, is coupled to




Fig. 1. Schematic representation of the Wood–Werkman cycle in *P. freudenreichii*. Reactions are directed toward propionate production, but are reversible.

oxidative phosphorylation and yields more ATP than in the other bacteria producing propionic acid.

Cells modulate the proportions of pyruvate either reduced to propionate, or oxidised to acetate and CO₂, to maintain the redox balance. The proportions of pyruvate directed to each pathway depend on the substrate used, the environmental conditions, and the properties of strains. For example, the oxidation of glucose and lactic acid leads to a molar ratio of propionate and acetate to ~2 (i.e. 2 moles of pyruvate are reduced to propionate while 1 mole of pyruvate is oxidised to acetate and CO₂). In contrast, the oxidation of glycerol leads to the formation of propionate only. *P. freudenreichii* is able to co-metabolise aspartate and other substrates like lactate, in a strain-dependant way. Aspartate is deaminated to fumarate that is further reduced to succinate, with a concomitant production of NAD and ATP. Strains using aspartate together with lactate convert less pyruvate to propionate and oxidised more pyruvate to acetate + CO₂, to maintain cell redox balance, compared to that metabolising lactate alone.

Acetate kinase gene disruption using integrational mutagenesis analysis on *P. acidipropionici* led to a decrease by 14% of acetate production from glucose and an increase by 13% of propionate yield (Suwannakham et al., 2006).

3.2. Respiration

P. freudenreichii is usually grown under anaerobic or microaerophilic conditions and described as an anaerobe. However, early studies also reported oxidative activity with free oxygen on a variety of substrates in propionibacteria (Vorobjeva, 1999). Accordingly, all the genes required for aerobic respiration were identified in *P. freudenreichii* genome: genes encoding NADH dehydrogenase, succinate dehydrogenase, cytochrome bd complex, ATPase and the complete pathway for heme synthesis (Falentin et al., 2010a). At high concentrations of oxygen, the synthesis of cytochrome, and thus the growth of *P. freudenreichii*, are inhibited. A switch from anaerobic to aerobic culture induces the consumption of the propionate produced in anaerobic conditions. Such a shift has been applied to improve the yield of vitamin B₁₂ production (Ye et al., 1999) and 1,4-dihydroxy-2-naphthoic acid production (Furuichi et al., 2007). Under anaerobic conditions, the electron acceptor in *P. freudenreichii* can be sulphate, fumarate, nitrate, menaquinone (vitamine K₂), or a pool of ferrous iron and humic acid in soil (Benz et al., 1998).

3.3. Vitamin and porphyrin synthesis

P. freudenreichii exhibits few nutritional requirements. It is able to synthesise all amino acids, and all but a few vitamins (Cummins and Johnson, 1992). Hence, *P. freudenreichii* can grow on chemically defined media containing a carbon and energy source, NH₄ as nitrogen source, minerals and vitamins (pantothenate, biotin, thiamine). *P. freudenreichii* synthesises in particular vitamin B₁₂ (cobalamin), a co-factor of methylmalonyl-CoA mutase, by the anaerobic pathway, regardless of the aerobic and anaerobic conditions of incubation (Iida et al., 2007; Roessner et al., 2002). Vitamin B₁₂ has been industrially produced for a long time and is the most complex vitamin synthesised by bacteria. More than 20 reactions are involved in the synthesis of this cobalt-containing molecule. The pathway of vitamin B₁₂ synthesis in *P. freudenreichii* has been thoroughly investigated (see reviews (Burgess et al., 2009; Murooka et al., 2005; Roessner et al., 2002)). This pathway was recently completed thanks to the complete genome sequence (Falentin et al., 2010a). Genetically modified strains of *P. freudenreichii* overproducing vitamins have been developed for vitamin B₁₂ (Murooka et al., 2005; Piao et al., 2004b), 5-aminolevulinic acid, the first intermediate in the synthesis of porphyrins (Kiatpapan and Murooka, 2001; Kiatpapan and Panbangred, 2008), porphyrin (Piao et al., 2004a), and riboflavin (vitamin B₂) (Burgess et al., 2009).

Genetically modified strains were obtained by the electroporation of *P. freudenreichii* IFO12426 with the *E. coli* shuttle vector pPK705 containing one to three genes of the respective pathways. Productions of vitamin B₂, vitamin B₁₂, and porphyrinogen were up to 2-fold, 1.9-fold and 33-fold greater, respectively, in overproducing strains, compared to the production in the wild strain (Burgess et al., 2006; Piao et al., 2004a; Piao et al., 2004b).

3.4. Production of exopolysaccharides

P. freudenreichii produces exopolysaccharides (EPS), including both hetero- and homopolysaccharides. Hitherto two primary EPS structures have been described in different strains: an heteropolysaccharide composed of D-glucose, D-mannose, and D-glucuronic acid in molar ratios of 2:2:1 (Dobrurowska et al., 2008) and a (1 → 3,1 → 2)-β-D-glucan homopolysaccharide (Deutsch et al., 2008; Nordmark et al., 2005). A polysaccharide of the latter type, tightly associated with the cell wall, was identified in 35% of *P. freudenreichii* strains (Deutsch et al., 2010). Interestingly, a single gene, *gtf*, is responsible for the synthesis of this surface polysaccharide, as revealed by inactivation experiments and heterologous expression in *Lactococcus lactis* (Deutsch et al., 2010). By using quantitative reverse transcription PCR, it was shown that the synthesis of this surface polysaccharide was correlated with the level of *gtf* expression; a minimum level of transcription is required for synthesis of a detectable amount of the polysaccharide. The presence of an IS element in the *gtf* promoting sequence may explain the strain-dependent expression of *gtf* (Deutsch et al., 2010).

3.5. Features allowing long-term survival

Cultures of dairy propionibacteria are known to remain viable for many months at room temperature (Cummins and Johnson, 1992). Indeed *P. freudenreichii* is able to survive and to remain active in various environments, including cheese, but also in the digest tract (Hervé et al., 2007). This feature can be explained by the ability of *P. freudenreichii* to accumulate various energy and carbon storage compounds, briefly described below.

P. freudenreichii accumulates polyphosphate (polyP) as energy storage. The presence of polyP was visualised using electron microscopy, as of electron-dense granules in cells (Clark et al., 1986; Vorobjeva, 1999). PolyP are linear polymers of tens or hundreds of orthophosphate residues linked by high-energy phosphoanhydride bonds. Their synthesis is catalysed by polyphosphate kinase (PPK) that transfers the terminal phosphate of ATP to polyP. PolyPs are known to enable microorganisms to respond to and tolerate stress. They have also been proposed as the modulators of quorum sensing and biofilm development, as shown by studies of *Pseudomonas* PPK mutants (Seufferheld et al., 2008). The accumulation of polyP also regulates the activation of the RNA polymerase RpoS and the synthesis of ppGpp, a major signalling component of the stringent response (Seufferheld et al., 2008). The enzymes using polyP instead of ATP catalyse reversible reactions, as for pyrophosphate phosphofructokinase in *P. freudenreichii*, involved both in glycolysis and neoglucogenesis (Meurice et al., 2004). *P. freudenreichii* also possesses a glucokinase using polyP or ATP as the phosphate donor (Phillips et al., 1993).

Glycogen is an α-1,4 linked, α-1,6 branched glucose polymer used for long-term energy storage in animal, fungi, bacteria, and archaeobacteria (Ball and Morell, 2003). A first experimental evidence of the ability of *P. freudenreichii* to synthesise glycogen was given by *in vivo* ¹³C NMR analysis of *P. freudenreichii* type strain cells grown in the presence of ¹³C glucose (Meurice, 2004). More recently, all the genes potentially related to glycogen metabolism were identified in the genome of the same strain (CIRM-BIA1^T=CIP103027^T) (Falentin et al., 2010a).

Trehalose is a non-reducing disaccharide found in many organisms, from bacteria to mammals, in which it exerts various functions related to stress adaptation due to its particular physical features. In bacteria, trehalose can be used as a carbon and energy source and is also accumulated as compatible solute (Arguelles, 2000). The detection of trehalose in propionibacteria has been early described. It is synthesised in all dairy propionibacteria species and is accumulated under various stress-inducing conditions (Cardoso et al., 2004; Rolin et al., 1995). The genes involved in trehalose synthesis in *P. freudenreichii* were recently described (Cardoso et al., 2007). The amounts of trehalose synthesised are highly strain-dependant in *P. freudenreichii* (Table 1).

P. freudenreichii has the ability to adapt to the various stresses that it encounters during fermentation processes and in variable environments. Industrial starters added to milk for cheese manufacture are generally proposed as lyophilisates, and have thus to adapt to osmotic stress. During Emmental cheese manufacture, *P. freudenreichii* has to withstand different successive stresses: heating over 50 °C, acidification of the curd to pH 5.2, osmotic stress due to the NaCl addition at the brining step, and low temperature (4 to 12 °C) during cheese ripening. Some strains also bear the acid- and bile-related stresses encountered in the digestive tract, which is a prerequisite of their use as probiotics. Interestingly, the cell machinery involved in stress adaptation in *P. freudenreichii* (Leverrier et al., 2004) was shown to be encoded by redundant multicopy genes (Falentin et al., 2010a). This is in agreement with the ability shown by *P. freudenreichii* to adapt efficiently to various conditions.

4. Recent advances in knowledge on *P. freudenreichii*

4.1. Evolution of published data about dairy propionibacteria

About 40 scientific publications about dairy propionibacteria have been published per year over the past two decades. Papers related to *P. freudenreichii* constitute more than 50% of these publications. Propionibacteria were mainly studied in the past for their particular metabolism and their application in cheese ripening, but the fields of research have evolved over the time and they are nowadays mainly

studied for their health properties. The main research fields covered over the past 3 years are depicted in Fig. 2. Only 10% of publications are currently related to cheese-related properties of propionibacteria. In contrast, publications related to human health have increased in number over the last decade and now constitute more than 40% of publications about dairy propionibacteria. They include studies about the various probiotic potentials, the synthesis of bifidogenic growth factor, vitamins, conjugated linoleic acid (CLA) and the potential of propionibacteria as protective cultures. Papers related to the use of propionibacteria as silage bacterial additives and as probiotics for animals account for about 20% of papers.

4.2. Role in aroma formation in cheese

P. freudenreichii is the main *Propionibacterium* species used in cheese manufacture. It is used in Swiss-type cheeses, in which it is responsible for the characteristic eyes due to the explosive production of CO₂ occurring when *P. freudenreichii* grows (for reviews see for example (Fröhlich-Wyder and Bachmann, 2004; Langsrud and Reinbold, 1973b; Thierry et al., 2010). Propionibacteria can also be used in the manufacture of various cheeses without eyes to enhance flavour formation (Ben Lawlor et al., 2003; Fernandez-Espla and Fox, 1998; Thierry et al., 2005a).

P. freudenreichii has a main role in the formation of Swiss-type cheese flavour. It produces flavour compounds from three main pathways: lactate and aspartate fermentation, amino acid catabolism, and fat hydrolysis (Thierry et al., 2004a). Many of these properties are strain-dependent. Table 1 gives some examples of phenotypic biodiversity related to the use of *P. freudenreichii* in cheese ripening. The ability to grow and propionic fermentation rate in cheese have long been known to be highly variable (Richoux and Kerjean, 1995). The ability to remain active in the presence of increased amounts of salt, in particular, depends on *P. freudenreichii* strains (Richoux et al., 1998).

Lactate fermentation by *P. freudenreichii* results in the formation of propionate, acetate (and CO₂), both acids being considered as flavour compounds in cheese. As described above, the metabolism of aspartate by propionibacteria during lactate fermentation results in

Table 1

Examples of phenotypic biodiversity related to the use of *P. freudenreichii* in cheese ripening and as a probiotic.

Target	Experimental conditions	Factor of variation	Reference
Trehalose accumulation	Chemically defined medium, pH 7.0 and 30 °C, cells harvested at mid-exponential growth phase	0 to 132 mg trehalose/g cell protein	(Cardoso et al., 2004)
Salt resistance in cheese	Mini Swiss-type cheeses, with 1, 2 and 3% salt in moisture (S/M)	Lactate consumed after 14 d ripening in the warm room: 12 to 13 g/kg at 1% S/M; 7 to 13 g/kg at 2% S/M	(Richoux et al., 1998)
Lysis	Potassium phosphate buffer, pH 6.2 for 24 h at 37 °C Potassium phosphate buffer, pH 7.0 for 48 h at 33 °C Pattern of autolytic enzymes after renaturing PAGE	Decrease of optical density: 24 to 96% Decrease of optical density: 24 to 62% Similar enzyme pattern	(Lemée et al., 1994) (Ostlie et al., 1995) (Ostlie et al., 2007)
Methylbutanoic acids	Emmental cheese, 8 strains 40 strains grown in lactate broth, pH 5.4, 21 g/L NaCl, at 24 °C	19 to 114 mg/kg cheese 6 to 57 mg/L culture supernatant	(Thierry et al., 2004c) (Dherbécourt et al., 2008b)
Free fatty acids (FFA) from lipolysis	Emmental cheese, 5 strains Lactate broth in the presence of an emulsion of milk fat	1.8 to 5.2 mg FFA/kg cheese Production of 0.8 to 3 mg FFA/kg fat	(Chamba and Perréard, 2002) (Dherbécourt et al., 2010b) for culture conditions; Thierry, unpublished data
Aspartase activity	Incubation of cell-free extract (12 strains) in the presence of aspartate – quantification of fumarate produced	<0.5 to 35 mmol/min/mg protein	(Turgay et al., 2010)
Growth at low temperature	Lactate broth, pH 5.5, incubated at 11 °C	<0.1 to 2.0 units of optical density (OD ₅₇₈) after 24 days	(Turgay et al., 2010)
Tolerance to acid and bile	Acid + bile stress	cfu decrease 0.5 to 4 log	(Lan et al., 2007a)
Propionate production	Medium designed to mimic the content of the human colon	0.4 to 1.4 g/L	(Lan et al., 2007a)
Surface polysaccharide	Agglutination tests (anti- <i>S. pneumoniae</i> ser 37 antiserum)	Positive/negative	(Deutsch et al., 2008; Deutsch et al., 2010)
IL-10 induction	Elisa on stimulated peripheral blood mononuclear cells	1500 to 5500 pg/mL	(Foligné et al., 2010)




Fig. 2. Fields of research covered by publications about dairy propionibacteria since 2007 (from search in ISI Web Of Science database). DHNA, 4-dihydroxy-2-naphtoic acid; CLA, conjugated linoleic acid; EPS, exopolysaccharides.

additional CO₂ production. The balance of fermentation products in *P. freudenreichii* has important consequences on cheese ripening. The intensity of aspartate metabolism in *P. freudenreichii* is highly strain-dependent. Strains with a high ability to metabolise aspartate exhibit a higher fermentation rate and produce a higher proportion of CO₂ per mole of lactate consumed (Wyder et al., 2001). They can also be associated with an undesirable late fermentation in Emmental cheese, resulting in the formation of slits and cracks. This opening defect can occur, for example, during the ripening of Swiss Emmental cheese, which lasts for several months at 10–12 °C (Fröhlich-Wyder and Bachmann, 2004). The specific aspartase activity (L-aspartate ammonia-lyase), compared in the cell-free extracts of 12 *P. freudenreichii* strains, varied from less than 0.5 to 35 mmol/min/mg protein (Turgay et al., 2010) (Table 1).

P. freudenreichii catabolises branched-chain amino acids to branched-chain volatile (flavour) compounds. The main branched-chain compounds produced by *P. freudenreichii* are 2-methylbutanoic acid and 3-methylbutanoic acid (actual isovaleric acid), which derive from isoleucine and leucine degradation, respectively (Thierry and Maillard, 2002). These two acids are important flavour compounds in many cheeses and are associated with typical “old cheese”, “sweaty socks” notes. *P. freudenreichii* is by far the main agent of the formation of methylbutanoic acids (comprising about 80% of 2-methylbutanoic acid and about 20% of 3-methylbutanoic acid in Swiss cheese) (Thierry et al., 2004b). The production of methylbutanoic acids is constitutive in *P. freudenreichii* and is related to the synthesis of long branched-chain fatty acids which constitute more than 80% of cell membrane fatty acids (Dherbécourt et al., 2008b). The synthesis of methylbutanoic acids occurs via transamination of branched-chain amino acids, leading to ketoacids that are further converted into acids by oxidative decarboxylation (Thierry and Maillard, 2002). Methylbutanoic acids are produced in broth cultures and in cheese by *P. freudenreichii*. The screening of *P. freudenreichii* strains ability to produce methylbutanoic acids gave similar results *in vitro* and in Swiss-type cheese (Thierry et al., 2004c). The concentrations of methylbutanoic acids produced in a modified lactate broth incubated at 24 °C varied by a factor of 8 within a pool of 40 strains (cf. Table 1). Four strains were distinguished by their significant higher ability to release methylbutanoic acids (>30 mg/L) (Dherbécourt et al., 2008b).

P. freudenreichii has also a prominent role in Emmental cheese lipolysis. It is responsible for almost the whole free fatty acids released from cheese fat during Swiss cheese ripening (Dherbécourt et al., 2010a). Free fatty acids are important flavour compounds in many cheeses, including Swiss cheese. The lipolytic activity of *P. freuden-*

reichii is strain-dependent (Chamba and Perréard, 2002) (Thierry, unpublished results). The most probable esterases potentially involved in cheese lipolysis were identified from the complete genome sequence of *P. freudenreichii* (Dherbécourt et al., 2008a). One of them was identified as the sole secreted esterase in *P. freudenreichii* by *in silico* and biochemical approaches. Transformed clones over-expressing this enzyme showed 5 to 8 times more lipolytic activity on milk fat than the wild-type strain mutant (Dherbécourt et al., 2010b). This result strongly indicates that this enzyme plays a major role in cheese lipolysis.

In contrast, *P. freudenreichii* has very low caseinolytic activity. It possesses diverse intracellular peptidases, including several enzymes specific of proline-containing peptides. However, *P. freudenreichii* has a limited role in secondary proteolysis of cheese, compared to lactic acid bacteria, because it does not lyse in cheese or only lately (Valence et al., 1998), even if the ability of strains to lyse *in vitro* in buffer varies over large ranges (Table 1) (Ostlie et al., 1995).

The production of flavour compounds occurs during *P. freudenreichii* growth in cheeses at about 24 °C, and continues during further cold storage of cheeses, while propionic fermentation is markedly slowed down at low temperatures. For example, about 60% of methylbutanoic acids from amino acid catabolism and 40% of fatty acids from lipolysis were produced during cold storage of experimental Swiss cheeses, while only 20% of propionic acid were produced during the same period (Thierry et al., 2005b). New insights are now made possible by the use of molecular methods to evaluate the activity of micro-organisms in cheese. It was recently shown that *P. freudenreichii* maintained metabolic activity up to the end of ripening, as shown by using quantitative reverse transcription PCR (RT-qPCR) specifically targeting different housekeeping genes (*16S*, *groL1*, *groL2* and *tuf*) (Falentin et al., 2010b). The ability to grow at low temperature is also highly strain-dependent (Table 1). This property, along with a high aspartase activity, would be important in the ability of *P. freudenreichii* strains added as cultures in Swiss cheese to maintain as the dominant population over long ripening periods of time (Turgay et al., 2010).

4.3. Probiotic properties of *P. freudenreichii*

P. freudenreichii and *P. acidipropionici* were considered for their potent probiotic effects on animals and on humans later than, for instance, lactobacilli and bifidobacteria, and scientific reports began in the 90s. Their probiotic potential was early reviewed in 1999 by A. Perez-Chaia et al., 1999 and more recently by Ouwehand, 2004 and Cousin et al., 2010. Briefly, dairy propionibacteria are considered for their beneficial modulation of the gut microbiota and of the corresponding metabolic activities. *P. freudenreichii* has also more recently been studied for its modulation of the gut function and physiology. We will focus here on the reports published during the last 5 years on *P. freudenreichii* and giving new mechanistic highlights on dairy propionibacteria probiotic effects with a potential application to the human health. Clinical studies only based on the use of complex probiotic mixtures including propionibacteria as probiotic are not covered by this paper, neither are probiotic applications in animals.

The bifidogenic effect of selected strains of *P. freudenreichii* was observed in independent studies performed on human volunteers (Bouglé et al., 1999; Hojo et al., 2002; Satomi et al., 1999). Volunteers consuming propionibacteria displayed enhanced colic bifidobacterial populations. The active compound was present in propionibacterial supernatants and identified as 1,4-dihydroxy-2-naphtoic acid (DHNA) (Isawa et al., 2002). This molecule is the penultimate intermediate in the biosynthesis pathway of vitamin K₂ (menaquinone). *P. freudenreichii* synthesises vitamin K₂ and the whole pathway was reconstructed from the genome data (Falentin et al., 2010a). DHNA is synthesised from 2-succinylbenzoyl-CoA by a naphtoate synthase and

then leaks out of propionibacteria without cell lysis. It has been proposed that DHNA and its derivative 2-amino-3-carboxy-1,4-naphthoquinone (ACNQ) serve as electron transfer mediator for NADP regeneration in bifidobacteria (Yamazaki et al., 1998; Yamazaki et al., 1999), thus favouring bifidobacterial growth. Besides the microbiota, Propionibacteria are also proposed to modulate metabolic activity in the favour of beneficial metabolites.

Production of beneficial conjugated linoleic acids (CLAs) is also documented in *P. freudenreichii*. Among the biological effects described for CLAs, the anticarcinogenic properties of rumenic acid, the cis-9,trans-11 stereoisomer of CLA, are highly promising. *P. freudenreichii* has been shown to convert free linoleic acid into rumenic acid (Rainio et al., 2002; Wang et al., 2007). The corresponding mechanisms of CLA formation by *P. freudenreichii* were identified by deuterium labelling and mass spectrometry (McIntosh et al., 2009). However, the genes encoding enzymes involved in this pathway are still not elucidated and the ability of *P. freudenreichii* to produce substantial amounts of rumenic acid *in situ* in fermented products or in the colic content remains largely to be studied. Several processes of enrichment of CLA, based on a lipid hydrolysis followed by a *P. freudenreichii*-catalysed isomerisation of free linoleic acid to CLA, have been developed in various products like oats (Vahvaselka et al., 2006) and by-products of plant oil processing (Vahvaselka and Laakso, 2010).

Beneficial metabolic activity is mainly limited by the ability of dairy propionibacteria to adapt stressing environments. *P. freudenreichii* can adapt to very unfavourable conditions encountered either during industrial processes or within the digestive tract (Leverrier et al., 2004; Suomalainen et al., 2008). The corresponding adaptation proteins were particularly investigated in *P. freudenreichii* and its genome revealed a remarkable redundancy in the cell machinery involved in stress perception, adaptation, detoxification and macromolecules repair (Falentin et al., 2010a). We have shown these proteins to be differentially expressed, depending on the strain (Anastasiou et al., 2006). Accordingly, the ability to adapt and maintain an active metabolism within the colon of human microbiota-associated rats and of human volunteers was shown to be highly strain dependent (Hervé et al., 2007; Lan et al., 2007a) (Table 1). This illustrates a notable phenotypic biodiversity among strains. In the colon, selected strains were showed to express the transcarboxylase operon specific of propionic fermentation, and to produce propionate. Propionate is known for its role in the modulation of the proliferation/apoptosis balance. Accordingly, the most promising strain selected in *in vivo* and *in vitro* screening was shown to reduce proliferation yet enhance apoptosis of colon epithelial cells in rats mutagenized with the carcinogen dimethylhydrazine (Lan et al., 2008). Interestingly, propionibacteria had no effect on these parameters in healthy rats. The corresponding pro-apoptotic mechanisms were well described *in vitro* on cultured human colon cancer cells and were shown to be enhanced by an acidic environment (Lan et al., 2007b). This pro-apoptotic property may prove useful in the context of colon cancer prevention and/or treatment. In the same context, one should notice that the ability of *P. freudenreichii* to bind the food born carcinogen aflatoxin B-1 *in vitro* (El-Nezami et al., 2000) was confirmed in a human study (El-Nezami et al., 2006). Human volunteers chronically exposed to this carcinogen showed reduced bioavailability and urinary excretion of aflatoxin, as a result of probiotic consumption. However, this probiotic containing both *Lactobacillus rhamnosus* and *P. freudenreichii*, the specific role of this last is difficult to specify.

Finally, immunomodulation by dairy propionibacteria was confirmed in humans, as volunteers consuming *P. freudenreichii* JS showed reduced serum levels of CRP, indicating an anti-inflammatory effect (Kekkonen et al., 2008). Indeed, immunomodulation was evidenced in a highly strain dependent manner in dairy propionibacteria (Foligné et al., 2010). Selected strains induced high levels of regulatory anti-inflammatory cytokines in human peripheral blood mononuclear cells (PBMCs) and were effective in two colitis models in

mice. The surface antigens of propionibacteria were shown to play a key role in such a modulation (Foligné et al., 2010). This is consistent with a previous study showing immunomodulatory properties of *P. freudenreichii* strain JS (Kekkonen et al., 2008).

The activity of *P. freudenreichii* against the pathogen bacterium *Helicobacter pylori* was also recently investigated. *In vitro*, *P. freudenreichii* reduced adhesion of *H. pylori* on intestinal cells and *H. pylori*-induced cell membrane leakage (Myllyluoma et al., 2008). Besides, it inhibited *H. pylori*-induced IL-8 and PGE2 release by intestinal cells. In clinical studies, the supplementation of a mixture of probiotics containing *P. freudenreichii* improved the tolerance to the treatment of *H. pylori* infection (Myllyluoma et al., 2005; Myllyluoma et al., 2007a) and led to a beneficial effect on gastric mucosa in *H. pylori*-infected patients (Myllyluoma et al., 2007b). Moreover, the bifidogenic growth stimulator DHNA produced by *P. freudenreichii* inhibited the growth of *H. pylori*, via the inhibition of cellular respiration (Nagata et al., 2010). Taken altogether, these studies suggest that *P. freudenreichii* may be useful to eradicate *H. pylori*.

4.4. Protective cultures

Some dairy propionibacteria can also synthesise anti-fungal and anti-yeast compounds and have a potential as food-protective cultures (Ho et al., 2009; Lind et al., 2007; Schwenninger et al., 2008; Tharmaraj and Shah, 2009). *P. freudenreichii* secretes bacteriocins. A recent review reports the literature on three antimicrobial peptides (PAMP antimicrobial peptide and propionins T1 and F) from propionibacteria, relating the genetics, biochemistry, biosynthesis, and biological activities of these compounds (Faye et al., 2010). In *P. jensenii*, several antimicrobial organic acids acting in synergy have recently been identified, including not only acetic, propionic, succinic and lactic acids, but also 2-pyrrolidone-5-carboxylic acid, 3-phenyllactic acid, hydroxyphenyl lactic acid 3-phenyllactic acid (Schwenninger et al., 2008).

5. Concluding remarks

As a conclusion, *P. freudenreichii* just entered the so-called post-genomic era and this will favour its use in various domains. Indeed, this bacterium has long attracted attention because of its peculiar properties and of its several useful applications. However, molecular tools and knowledge were lacking to deepen the general understanding of its action. The newly available genome sequence, together with the developed molecular tools, is opening new perspectives in the context of its use. The molecular mechanisms involved in key pathways for adaptation, cheese ripening, bio-preservation and probiotic effects can now be elucidated. This will in turn allow selection of the most appropriate strain for each application, taking advantage of *P. freudenreichii* remarkable biodiversity.

References

- Anastasiou, R., Leverrier, P., Krestas, I., Rouault, A., Kalantzopoulos, G., Boyaval, P., Tsakalidou, E., Jan, G., 2006. Changes in protein synthesis during thermal adaptation of *Propionibacterium freudenreichii* subsp. *shermanii*. International Journal of Food Microbiology 108, 301–314.
- Arguelles, J.C., 2000. Physiological roles of trehalose in bacteria and yeasts: a comparative analysis. Archives of Microbiology 174, 217–224.
- Ball, S.G., Morell, M.K., 2003. From bacterial glycogen to starch: understanding the biogenesis of the plant starch granule. Annual Review of Plant Biology 54, 207–233.
- Ben Lawlor, J., Delahunty, C.M., Wilkinson, M.G., Sheehan, J., 2003. Swiss-type and Swiss-Cheddar hybrid-type cheeses: effects of manufacture on sensory character and relationships between the sensory attributes and volatile compounds and gross compositional constituents. International Journal of Dairy Technology 56, 39–51.
- Benz, M., Schink, B., Brune, A., 1998. Humic acid reduction by *Propionibacterium freudenreichii* and other fermenting bacteria. Applied and Environmental Microbiology 64, 4507–4512.
- Bernard, K.A., Shuttleworth, L., Munro, C., Forbes-Faulkner, J.C., Pitt, D., Norton, J.H., Thomas, A.D., 2002. *Propionibacterium australiense* sp. nov. derived from granulomatous bovine lesions. Anaerobe 8, 41–47.

- Bouglé, D., Roland, N., Lebeurier, F., Arhan, P., 1999. Effect of propionibacteria supplementation on fecal bifidobacteria and segmental colonic transit time in healthy human subjects. *Scandinavian Journal of Gastroenterology* 2, 144–148.
- Burgess, C.M., Smid, E.J., Rutten, G., van, S.D., 2006. A general method for selection of riboflavin-overproducing food grade micro-organisms. *Microbial Cell Factories* 5, 24.
- Burgess, C.M., Smid, E.J., Van Sinderen, D., 2009. Bacterial vitamin B₂, B₁₁ and B₁₂ overproduction: an overview. *International Journal of Food Microbiology* 133, 1–7.
- Cardoso, F.S., Gaspar, P., Hugenholtz, J., Ramos, A., Santos, H., 2004. Enhancement of trehalose production in dairy propionibacteria through manipulation of environmental conditions. *International Journal of Food Microbiology* 91, 195–204.
- Cardoso, F.S., Castro, R.F., Borges, N., Santos, H., 2007. Biochemical and genetic characterization of the pathways for trehalose metabolism in *Propionibacterium freudenreichii*, and their role in stress response. *Microbiology* 153, 270–280.
- Chamba, J.F., Perréard, E., 2002. Contribution of propionibacteria to lipolysis of Emmental cheese. *Le Lait* 82, 33–44.
- Clark, J.E., Beegen, H., Wood, H.G., 1986. Isolation of intact chains of polyphosphate from *Propionibacterium shermanii* grown on glucose or lactate. *Journal of Bacteriology* 168, 1212–1219.
- Cousin, F., Mater, D.D.G., Foligné, B., Jan, G., 2010. Dairy propionibacteria as human probiotics: a review of recent evidence. *Dairy Science & Technology*. doi:10.1051/dst/2010032-.
- Crow, V.L., 1987. Citrate cycle intermediates in the metabolism of aspartate and lactate by *Propionibacterium freudenreichii* subsp. *shermanii*. *Applied and Environmental Microbiology* 53, 2600–2602.
- Cummins, C.S., Johnson, J.L., 1986. Genus I. *Propionibacterium* Orla-Jensen 1909. In: Sneath, P.H.A., Mair, N.S., Sharpe, M.E., Holt, J.G. (Eds.), *Bergey's Manual of Systematic Bacteriology*. Williams & Wilkins, Baltimore, pp. 1346–1353.
- Cummins, C.S., Johnson, J.L., 1992. The genus *Propionibacterium*. In: Balows, E., Truper, H.G., Dworkin, M., Harder, W., Schleifer, K.H. (Eds.), *The Prokaryotes*. Springer Verlag.
- Dalmasso, M., Nicolas, P., Falentin, H., Valence, F., Tanskanen, J., Jatila, H., Salusjärvi, T., Thierry, A., 2011. Multilocus sequence typing of *Propionibacterium freudenreichii*. *International Journal of Food Microbiology* 145, 113–120.
- de Carvalho, A.F., Gautier, M., Grimont, F., 1994. Identification of dairy *Propionibacterium* species by rRNA gene restriction patterns. *Research in Microbiology* 145, 667–676.
- Deutsch, S.M., Falentin, H., Dols-Lafargue, M., LaPointe, G., Roy, D., 2008. Capsular exopolysaccharide biosynthesis gene of *Propionibacterium freudenreichii* subsp. *shermanii*. *International Journal of Food Microbiology* 125, 252–258.
- Deutsch, S.M., Le Bivic, P., Herve, C., Madec, M.N., LaPointe, G., Jan, G., Le Loir, Y., Falentin, H., 2010. Correlation of the capsular phenotype in *Propionibacterium freudenreichii* with the level of expression of *gtf*, a unique polysaccharide synthase-encoding gene. *Applied and Environmental Microbiology* 76, 2740–2746.
- Dherbécourt, J., Falentin, H., Canaan, S., Thierry, A., 2008a. A genomic search approach to identify esterases in *Propionibacterium freudenreichii* involved in the formation of flavour in Emmental cheese. *Microbial Cell Factories* 7.
- Dherbécourt, J., Maillard, M.B., Catheline, D., Thierry, A., 2008b. Production of branched-chain aroma compounds by *Propionibacterium freudenreichii*: links with the biosynthesis of membrane fatty acids. *Journal of Applied Microbiology* 105, 977–985.
- Dherbécourt, J., Bourlieu, C., Maillard, M.B., Aubert-Frogerais, L., Richoux, R., Thierry, A., 2010a. Time course and specificity of lipolysis in Swiss cheese. *Journal of Agricultural and Food Chemistry* 58, 11732–11739.
- Dherbécourt, J., Falentin, H., Jardin, J., Maillard, M.B., Baglinière, F., Barloy-Hubler, F., Thierry, A., 2010b. Identification of a secreted lipolytic esterase in *Propionibacterium freudenreichii*, a ripening process bacterium involved in Emmental cheese lipolysis. *Applied and Environmental Microbiology* 76, 1181–1188.
- Dobrurowska, J.M., Gerwig, G.J., Babuchowski, A., Kamerling, J.P., 2008. Structural studies on exopolysaccharides produced by three different propionibacteria strains. *Carbohydrate Research* 343, 726–745.
- Dorner, W., 1939. Recherches sur les bactéries propioniques. *Le Lait* 19, 897–918.
- Downes, J., Wade, W.G., 2009. *Propionibacterium acidifaciens* sp. nov., isolated from the human mouth. *International Journal of Systematic and Evolutionary Microbiology* 59, 2778–2781.
- EFSA, 2008. Technical guidance update of the criteria used in the assessment of bacterial resistance to antibiotics of human or veterinary importance. *The EFSA Journal* 732, 1–15.
- EFSA, 2009. Scientific opinion of the panel on biological hazards on the maintenance of the list of QPS microorganisms intentionally added to food or feed. *The EFSA Journal* 7, 1–93.
- El-Nezami, H., Mykkanen, H., Kankaanpää, P., Salminen, S., Ahokas, J., 2000. Ability of *Lactobacillus* and *Propionibacterium* strains to remove aflatoxin B₁ from the chicken duodenum. *J. Food* 63, 549–552.
- El-Nezami, H.S., Polychronaki, N.N., Ma, J., Zhu, H., Ling, W., Salminen, E.K., Juvonen, R.O., Salminen, S.J., Poussa, T., Mykkanen, H.M., 2006. Probiotic supplementation reduces a biomarker for increased risk of liver cancer in young men from Southern China. *The American Journal of Clinical Nutrition* 83, 1199–1203.
- Falentin, H., Deutsch, S.M., Jan, G., Loux, V., Thierry, A., Parayre, S., Maillard, M.B., Dherbécourt, J., Cousin, F.J., Jardin, J., Siguier, P., Couloux, A., Barbe, V., Vacherie, B., Wincker, P., Gibrat, J.F., Gaillardin, C., Lortal, S., 2010a. The complete genome of *Propionibacterium freudenreichii* CIRM-BIA1, a hardy actinobacterium with food and probiotic applications. *PLoS One* 5, e11748.
- Falentin, H., Postollec, F., Parayre, S., Henaff, N., Le, B.P., Richoux, R., Thierry, A., Sohier, D., 2010b. Specific Metabolic Activity of Ripening Bacteria Quantified by Real-time Reverse Transcription PCR throughout Emmental Cheese Manufacture. *International Journal of Food Microbiology* 144, 10–19.
- Faye, T., Holo, H., Langsrud, T., Nes, I.F., Brede, D.A., 2010. The Unconventional Antimicrobial Peptides of the Classical Propionibacteria. *Applied Microbiology and Biotechnology* 89, 549–554.
- Fernandez-Espla, M.D., Fox, P.F., 1998. Effect of adding *Propionibacterium shermanii* NCD0 853 or *Lactobacillus casei* ssp. *casei* IPPL 731 on proteolysis and flavor development of Cheddar cheese. *Journal of Agricultural and Food Chemistry* 46, 1228–1234.
- Fessler, D. S., 1997. Characterisation of propionibacteria in Swiss raw milk by biochemical and molecular-biological methods. PhD thesis, Swiss Federal Institute of Technology, Zurich, Suisse.
- Foligné, B., Deutsch, S.M., Breton, J., Cousin, F.J., Dewulf, J., Samson, M., Pot, B., Jan, G., 2010. Promising immunomodulatory effects of selected strains of dairy propionibacteria evidenced *in vitro* and *in vivo*. *Applied and Environmental Microbiology* 76, 8259–8264.
- Fröhlich-Wyder, M.T., Bachmann, H.P., 2004. Cheeses with propionic acid fermentation. In: Fox, P.F., McSweeney, P.L.H., Cogan, T.M., Guinee, T.P. (Eds.), *Cheese. Chemistry, Physics and Microbiology*. Elsevier, London, pp. 141–156.
- Furuichi, K., Katakura, Y., Ninomiya, K., Shioya, S., 2007. Enhancement of 1,4-dihydroxy-2-naphthoic acid production by *Propionibacterium freudenreichii* ET-3 fed-batch culture. *Applied and Environmental Microbiology* 73, 3137–3143.
- Hervé, C., Fondrevez, M., Chéron, A., Barloy-Hubler, F., Jan, G., 2007. Transcarboxylase mRNA: a marker which evidences *P. freudenreichii* survival and metabolic activity during its transit in the human gut. *International Journal of Food Microbiology* 113, 303–314.
- Hettinga, D.H., Reinbold, G.W., 1972. The propionic-acid bacteria – a review II metabolism. *Journal of Milk and Food Technology* 35, 358–372.
- Ho, P.H., Luo, J.B., Adams, M.C., 2009. *Lactobacilli* and dairy *Propionibacterium* with potential as biopreservatives against food fungi and yeast contamination. *Applied Biochemistry and Microbiology* 45, 414–418.
- Hojo, K., Yoda, N., Tsuchita, H., Ohtsu, T., Seki, K., Taketomo, N., Murayama, T., Iino, H., 2002. Effect of ingested culture of *Propionibacterium freudenreichii* ET-3 on fecal microflora and stool frequency in healthy females. *Bioscience and Microflora* 21, 115–120.
- Houwen, F.P., Dijkema, C., Stams, A.J.M., Zehnder, A.J.B., 1991. Propionate metabolism in anaerobic bacteria; determination of carboxylation reactions with ¹³C-NMR spectroscopy. *Biochimica et Biophysica Acta (BBA) – Bioenergetics* 1056, 126–132.
- Iida, K., Ohtaka, K., Kajiwara, M., 2007. Mechanism of the ring contraction process in vitamin B-12 biosynthesis by the anaerobe *Propionibacterium shermanii* under aerobic conditions. *The FEBS Journal* 274, 3475–3481.
- Isawa, K., Hojo, K., Yoda, N., Kamiyama, T., Makino, S., Saito, M., Sugano, H., Mizoguchi, C., Kurama, S., Shibasaki, M., Endo, N., Sato, Y., 2002. Isolation and identification of a new bifidogenic growth stimulator produced by *Propionibacterium freudenreichii* ET-3. *Bioscience, Biotechnology, and Biochemistry* 66, 679–681.
- Kekkonen, R.A., Lummela, N., Karjalainen, H., Latvala, S., Tynkkynen, S., Jarvenpää, S., Kautiainen, H., Julkunen, I., Vapaatalo, H., Korpela, R., 2008. Probiotic intervention has strain-specific anti-inflammatory effects in healthy adults. *World Journal of Gastroenterology* 14, 2029–2036.
- Kiatpapan, P., Murooka, Y., 2001. Construction of an expression vector for propionibacteria and its use in production of 5-aminolevulinic acid by *Propionibacterium freudenreichii*. *Applied Microbiology and Biotechnology* 56, 144–149.
- Kiatpapan, P., Panbangred, W., 2008. Production of 5-aminolevulinic acid by propionibacteria. *The FEBS Journal* 275, 407–407.
- Koussemon, M., Combet-Blanc, Y., Patel, B.K., Cayol, J.L., Thomas, P., Garcia, J.L., Ollivier, B., 2001. *Propionibacterium microaerophilum* sp. nov., a microaerophilic bacterium isolated from olive mill wastewater. *International Journal of Systematic and Evolutionary Microbiology* 51, 1373–1382.
- Kusano, K., Yamada, H., Niwal, M., Yamasato, K., 1997. *Propionibacterium cyclohexanicum* sp. nov., a new acid-tolerant omega-cyclohexyl fatty acid-containing *Propionibacterium* isolated from spoiled orange juice. *International Journal of Systematic Bacteriology* 47, 825–831.
- Lan, A., Bruneau, A., Philippe, C., Rochet, V., Rouault, A., Hervé, C., Roland, N., Rabot, S., Jan, G., 2007a. Survival and metabolic activity of selected strains of *Propionibacterium freudenreichii* in the gastrointestinal tract of human microbiota-associated rats. *The British Journal of Nutrition* 97, 714–724.
- Lan, A., Lagadic-Gossmann, D., Lemaire, C., Brenner, C., Jan, G., 2007b. Acidic extracellular pH shifts colorectal cancer cell death from apoptosis to necrosis upon exposure to propionate and acetate major end-products of the human probiotic propionibacteria. *Apoptosis* 12, 573–591.
- Lan, A., Bruneau, A., Bensaada, M., Philippe, C., Bellaud, P., Rabot, S., Jan, G., 2008. Increased induction of apoptosis by *Propionibacterium freudenreichii* TL133 in colonic mucosal crypts of human microbiota-associated rats treated with 1,2-dimethylhydrazine. *The British Journal of Nutrition* 1–9.
- Langsrud, T., Reinbold, G.W., 1973a. Flavor development and microbiology of Swiss cheese – a review. II. Starters, manufacturing processes and procedures. *Journal of Milk and Food Technology* 36, 531–542.
- Langsrud, T., Reinbold, G.W., 1973b. Flavor development and microbiology of Swiss cheese – a review III. Ripening and flavor production. *Journal of Milk and Food Technology* 36, 593–609.
- Lemée, R., Rouault, A., Guezenc, S., Lortal, S., 1994. Autolysis of 57 strains of dairy propionibacteria. *Le Lait* 74, 241–251.
- Leuschner, R.G.K., Robinson, T.P., Hugas, M., Cocconcelli, P.S., Richard-Forget, F., Klein, G., Licht, T.R., Nguyen-the, C., Querol, A., Richardson, M., Suarez, J.E., Thrane, U., Vlak, J.M., von, W.A., 2010. Qualified presumption of safety (QPS): a generic risk assessment approach for biological agents notified to the European Food Safety Authority (EFSA). *Trends in Food Science & Technology* 21, 21.
- Leverrier, P., Vissers, J.P.C., Rouault, A., Boyaval, P., Jan, G., 2004. Mass spectrometry proteomic analysis of stress adaptation reveals both common and distinct response pathways in *Propionibacterium freudenreichii*. *Archives of Microbiology* 181, 215–230.

- Lind, H., Sjogren, J., Gohil, S., Kenne, L., Schnurer, J., Broberg, A., 2007. Antifungal compounds from cultures of dairy propionibacteria type strains. *FEMS Microbiology Letters* 271, 310–315.
- McIntosh, F.M., Shingfield, K.J., Devillard, E., Russell, W.R., Wallace, R.J., 2009. Mechanism of conjugated linoleic acid and vaccenic acid formation in human faecal suspensions and pure cultures of intestinal bacteria. *Microbiology-Sgm* 155, 285–294.
- Meile, L., Le Blay, G., Thierry, A., 2008. Contribution to the safety assessment of technological microflora found in fermented dairy products Part IX. *Propionibacterium* and *Bifidobacterium*. *International Journal of Food Microbiology* 126, 316–320.
- Meurice, G., 2004. Reconstruction *in silico* de voies métaboliques : application aux voies glycolytiques de *Propionibacterium freudenreichii* subsp. *shermanii*. PhD thesis, Ecole Nationale Supérieure Agronomique de Rennes, Rennes, France.
- Meurice, G., Deborde, C., Jacob, D., Falentin, H., Boyaval, P., Dimova, D., 2004. In silico exploration of the fructose-6-phosphate phosphorylation step in glycolysis: genomic evidence of the coexistence of an atypical ATP-dependent along with a PPI-dependent phosphofruktokinase in *Propionibacterium freudenreichii* subsp. *shermanii*. *In Silico Biology* 4, 0043.
- Mogensen, G., Salminen, S., O'Brien, J., Ouwehand, A., Holzapfel, W., Shortt, C., Fonden, R., Miller, G.D., Donohue, D., Playne, M., Crittenden, R., Salvadori, B.B., Zink, R., 2002. Inventory of microorganisms with a documented history of use in food. *Bulletin of the International Dairy Federation* 10–19.
- Moore, W.E.C., Holdeman, L.V., 1986. *Propionibacterium*. In: Buchanan, R.E., Gibbons, R.E. (Eds.), *Bergey's Manual of Determinative Bacteriology*. Williams & Wilkins, Baltimore, pp. 633–641.
- Murooka, Y., Piao, Y., Kiatpapan, P., Yamashita, M., 2005. Production of tetrapyrrole compounds and vitamin B₁₂ using genetically engineering of *Propionibacterium freudenreichii* An overview. *Le Lait* 85, 9–22.
- Myllyluoma, E., Veijola, L., Ahlroos, T., Tynkkynen, S., Kankuri, E., Vapaatalo, H., Rautelin, H., Korpela, R., 2005. Probiotic supplementation improves tolerance to *Helicobacter pylori* eradication therapy – a placebo-controlled, double-blind randomized pilot study. *Alimentary Pharmacology & Therapeutics* 21, 1263–1272.
- Myllyluoma, E., Ahlroos, T., Veijola, L., Rautelin, H., Tynkkynen, S., Korpela, R., 2007a. Effects of anti-*Helicobacter pylori* treatment and probiotic supplementation on intestinal microbiota. *International Journal of Antimicrobial Agents* 29, 66–72.
- Myllyluoma, E., Kajander, K., Mikkola, H., Kyrönpalo, S., Rasmussen, M., Kankuri, E., Sipponen, P., Vapaatalo, H., Korpela, R., 2007b. Probiotic intervention decreases serum gastrin-17 in *Helicobacter pylori* infection. *Digestive and Liver Disease* 39, 516–523.
- Myllyluoma, E., Ahonen, A.M., Korpela, R., Vapaatalo, H., Kankuri, E., 2008. Effects of multispecies probiotic combination on *Helicobacter pylori* infection *in vitro*. *Clinical and Vaccine Immunology* 15, 1472–1482.
- Nagata, K., Inatsu, S., Tanaka, M., Sato, H., Kouya, T., Taniguchi, M., Fukuda, Y., 2010. The bifidogenic growth stimulator inhibits the growth and respiration of *Helicobacter pylori*. *Helicobacter* 15, 422–429.
- Nord, C.E., Oprica, C., 2006. Antibiotic resistance in *Propionibacterium acnes* microbiological and clinical aspects. *Anaerobe* 12, 207–210.
- Nordmark, E.L., Yang, Z.N., Huttunen, E., Widmalm, G., 2005. Structural studies of the exopolysaccharide produced by *Propionibacterium freudenreichii* ssp. *shermanii* JS. *Biomacromolecules* 6, 521–523.
- Ostlie, H., Vegarud, G., Langsrud, T., 1995. Autolysis of dairy propionibacteria in buffer systems. *Journal of Dairy Science* 78, 2315–2325.
- Ostlie, H.M., Vegarud, G., Langsrud, T., 2007. Autolysis of propionibacteria: detection of autolytic enzymes by renaturing SDS-PAGE and additional buffer studies 2160. *International Journal of Food Microbiology* 117, 167–174.
- Ouwehand, A.C., 2004. The probiotic potential of propionibacteria. In: Salminen, S., Ouwehand, A.C., Von Wright, A. (Eds.), *Lactic Acid Bacteria: Microbiology and Functional Aspects*. Marcel Dekker, Inc., New York, pp. 169–174.
- Perez-Chaia, A., Zarate, G., Oliver, G., 1999. The probiotic properties of propionibacteria. *Le Lait* 79, 175–185.
- Phillips, N.F.B., Horn, P.J., Wood, H.G., 1993. The polyphosphate-dependent and ATP-dependent glucokinase from *Propionibacterium shermanii* – both activities are catalyzed by the same protein. *Archives of Biochemistry and Biophysics* 300, 309–319.
- Piao, Y., Kiatpapan, P., Yamashita, M., Murooka, Y., 2004a. Effects of expression of hemA and hemB genes on production of porphyrin in *Propionibacterium freudenreichii*. *Applied Microbiology and Biotechnology* 70, 7561–7566.
- Piao, Y., Yamashita, M., Kawarachi, N., Asegawa, R., Ono, H., Murooka, Y., 2004b. Production of vitamin B12 in genetically engineered *Propionibacterium freudenreichii*. *Journal of Bioscience and Bioengineering* 98, 167–173.
- Piveteau, P., 1999. Metabolism of lactate and sugars by dairy propionibacteria: a review. *Le Lait* 79, 23–41.
- Rainio, A., Vahvaselka, M., Suomalainen, T., Laakso, S., 2002. Production of conjugated linoleic acid by *Propionibacterium freudenreichii* ssp. *shermanii*. *Le Lait* 82, 91–101.
- Richoux, R., Kerjean, J.R., 1995. Caractérisation technologique de souches pures de bactéries propioniques : test en minifabrication de fromages à pâte cuite. *Le Lait* 75, 45–59.
- Richoux, R., Favre, E., Kerjean, J.R., 1998. Effet de la teneur en NaCl sur la fermentation du lactate par *Propionibacterium freudenreichii* dans des minifromages à pâte cuite. *Le Lait* 78, 319–331.
- Roessner, C.A., Huang, K.X., Warren, M.J., Raux, E., Scott, A.I., 2002. Isolation and characterization of 14 additional genes specifying the anaerobic biosynthesis of cobalamin (vitamin B12) in *Propionibacterium freudenreichii* (P. *shermanii*). *Microbiology* 148, 1845–1853.
- Rolin, D.B., Girard, F., De Certaines, J.D., Boyaval, P., 1995. ¹³C-NMR study of lactate metabolism in *Propionibacterium freudenreichii* subsp. *shermanii*. *Applied Microbiology and Biotechnology* 44, 210–217.
- Satomi, K., Kurihara, H., Isawa, K., Mori, H., Kaneco, T., 1999. Effects of culture-powder of *Propionibacterium freudenreichii* ET-3 on fecal microflora of normal adults. *Bioscience, Biotechnology, and Biochemistry* 18, 27–30.
- Schwenninger, S.M., Lacroix, C., Truttmann, S., Jans, C., Spornli, C., Bigler, L., Meile, L., 2008. Characterization of low-molecular-weight antiyeast metabolites produced by a food-protective *Lactobacillus-Propionibacterium* coculture. *Journal of Food Protection* 71, 2481–2487.
- Seufferheld, M.J., Alvarez, H.M., Farias, M.E., 2008. Role of polyphosphates in microbial adaptation to extreme environments. *Applied and Environmental Microbiology* 74, 5867–5874.
- Stackebrandt, E., Rainey, A., Ward-Rainey, N.L., 1997. Proposal for a new hierarchic classification system, *Actinobacteria* classis nov. *International Journal of Systematic Bacteriology* 47, 479–491.
- Steffen, C., Eberhard, P., Bosset, J.O., Rüegg, M., 1993. Swiss-type varieties. In: Fox, P.F. (Ed.), *Cheese: Chemistry, Physics and Microbiology*. Elsevier Applied Science Publishers, London, UK, pp. 83–110.
- Suomalainen, T., Sigvart-Mattila, P., Matto, J., Tynkkynen, S., 2008. In vitro and in vivo gastrointestinal survival, antibiotic susceptibility and genetic identification of *Propionibacterium freudenreichii* ssp. *shermanii* JS. *International Dairy Journal* 18, 271–278.
- Suwannakham, S., Huang, Y., Yang, S.T., 2006. Construction and characterization of ack knock-out mutants of *Propionibacterium acidipropionici* for enhanced propionic acid fermentation. *Biotechnology and Bioengineering* 94, 94.
- Tharmaraj, N., Shah, N.P., 2009. Antimicrobial effects of probiotic bacteria against selected species of yeasts and moulds in cheese-based dips. *International Journal of Food Science and Technology* 44, 1916–1926.
- Thierry, A., Maillard, M.B., 2002. Production of cheese flavour compounds derived from amino acid catabolism by *Propionibacterium freudenreichii*. *Le Lait* 82, 17–32.
- Thierry, A., Maillard, M.B., Hervé, C., Richoux, R., Lortal, S., 2004a. Varied volatile compounds are produced by *Propionibacterium freudenreichii* in Emmentaler cheese. *Food Chemistry* 87, 439–446.
- Thierry, A., Richoux, R., Kerjean, J.R., 2004b. Isovaleric acid is mainly produced by *Propionibacterium freudenreichii* in Swiss cheese. *International Dairy Journal* 14, 801–807.
- Thierry, A., Richoux, R., Kerjean, J.R., Lortal, S., 2004c. A simple screening method for isovaleric acid production by *Propionibacterium freudenreichii* in Swiss cheese. *International Dairy Journal* 14, 697–700.
- Thierry, A., Maillard, M.B., Bonnarme, P., Roussel, E., 2005a. The addition of *Propionibacterium freudenreichii* to Raclette cheese induces biochemical changes and enhances flavour development. *Journal of Agricultural and Food Chemistry* 53, 4157–4165.
- Thierry, A., Maillard, M.B., Richoux, R., Kerjean, J.R., Lortal, S., 2005b. *Propionibacterium freudenreichii* strains quantitatively affect production of volatile compounds in Swiss cheese. *Le Lait* 85, 57–74.
- Thierry, A., Berthier, F., Gagnaire, V., Kerjean, J.R., Lopez, C., Noël, Y., 2010. Eye formation and Swiss-type cheeses. In: Law, B.A., Tamime, A.Y. (Eds.), *Technology of Cheesemaking*. Wiley-Blackwell, pp. 360–383.
- Thierry, A., Falentin, H., Deutsch, S.M., Jan, G., 2011. Bacteria, beneficial: *Propionibacterium* spp. In: Fuquay, J.W., Fox, P.F., McSweeney, P. (Eds.), *Encyclopedia of Dairy Science*. Elsevier, pp. 403–411.
- Turgay, M., Irmiler, S., Isolini, D., Wagner, E., Berthoud, H., Amrein, R., Fröhlich-Wyder, M.T., Wechsler, D., 2010. Characterization of wild-type and culture strains of propionic acid bacteria from raw milk and Emmentaler PDO cheese. Poster. 3rd International Symposium on Propionibacteria and Bifidobacteria, June 1–4 Oviedo, Spain.
- Vahvaselka, M., Laakso, S., 2010. Production of cis-9, trans-11-conjugated linoleic acid in camelina meal and okara by an oat-assisted microbial process. *Journal of Agricultural and Food Chemistry* 58, 2479–2482.
- Vahvaselka, M., Lehtinen, P., Laakso, S., 2006. Microbially safe utilization of non-inactivated oats (*Avena sativa* L.) for production of conjugated linoleic acid. *Journal of Agricultural and Food Chemistry* 54, 963–967.
- Valence, F., Richoux, R., Thierry, A., Palva, A., Lortal, S., 1998. Autolysis of *Lactobacillus helveticus* and *Propionibacterium freudenreichii* in Swiss cheeses: first evidence by using species-specific lysis markers. *The Journal of Dairy Research* 65, 609–620.
- Van Luijk, N., Stierli, M.P., Schwenninger, S.M., Herve, C., Dasen, G., Jore, J.P.M., Pouwels, P.H., van der Werf, M.J., Teuber, M., Meile, L., 2002. Genetics and molecular biology of propionibacteria. *Le Lait* 82, 45–57.
- Ventura, M., Canchaya, C., Tauch, A., Chandra, G., Fitzgerald, G.F., Chater, K.F., Van Sinderen, D., 2007. Genomics of Actinobacteria: tracing the evolutionary history of an ancient phylura. *Microbiology and Molecular Biology Reviews* 71, 495–548.
- von Freudenreich, E., Orla-Jensen, O., 1906. Ueber die in Emmentalerkäse stattfindende Propionsäure-gärung. *Zentralbl. Bakteriol.* 17, 529–546.
- Vorobjeva, L.L., 1999. *Propionibacteria*. Dordrecht, Kluwer Academic, pp. 1–291.
- Wang, L.M., Lv, J.P., Chu, Z.Q., Cui, Y.Y., Ren, X.H., 2007. Production of conjugated linoleic acid by *Propionibacterium freudenreichii*. *Food Chemistry* 103, 313–318.
- Wood, H.G., 1981. Metabolic cycles in the fermentation by propionic acid bacteria. *Current Topics in Cellular Regulation* 18, 255–287.
- Wyder, M.T., Bosset, J.O., Casey, M.G., Isolini, D., Sollberger, H., 2001. Influence of two different propionibacterial cultures on the characteristics of Swiss-type cheese with regard to aspartate metabolism. *Milchwissenschaften* 56, 78–81.
- Yamazaki, S., Kano, K., Ikeda, T., Isawa, K., Kaneko, T., 1998. Mechanistic study on the roles of a bifidogenic growth stimulator based on physicochemical characterization. *Biochimica et Biophysica Acta-General Subjects* 1425, 516–526.
- Yamazaki, S., Kano, K., Ikeda, T., Isawa, K., Kaneko, T., 1999. Role of 2-amino-3-carboxy-1,4-naphthoquinone, a strong growth stimulator for bifidobacteria, as an electron transfer mediator for NAD(P)(+) regeneration in *Bifidobacterium longum*. *Biochimica et Biophysica Acta-General Subjects* 1428, 241–250.
- Ye, K.M., Shijo, M., Miyano, K., Shimizu, K., 1999. Metabolic pathway of *Propionibacterium* growing with oxygen: enzymes, C-13 NMR analysis, and its application for vitamin B-12 production with periodic fermentation. *Biotechnology Progress* 15, 201–207.